



GLOBAL JOURNAL OF ADVANCED RESEARCH
(Scholarly Peer Review Publishing System)

RELATIONSHIP BETWEEN SOME CLIMATIC FACTORS AND SOIL CARBON DIOXIDE EMISSION

M. A. Youssef

Department of Soil and Water Science,
Fac. Agric., Al-Azhar Univ., Assuit,
Egypt.

I. I. Sadek

Central Laboratory for Agricultural Climate (CLAC),
Agricultural Research Center, Giza,
Egypt.

ABSTRACT

As the atmospheric CO₂ concentration continues to increase, more attention is being focused on the soil as a possible sink for atmospheric CO₂. Fertilizer application to soil can play a vital role in influencing the losses of soil carbon by CO₂ emission from the soil. This study was conducted to examine effect of different fertilization types on soil CO₂ emissions. Emission of CO₂ from wheat cultivated soil and fertilized with five different types of both organic and bio fertilizers were measured during seasons of 2011/2012 and 2012/2013, comparing with the emission in case of using chemical fertilizers at experimental farm and laboratory of Soils and Water Sci. Dept., Fac. of Agric., AL-Azhar Univ., Assiut, Egypt. Tested fertilizers were 150 L/fed., compost tea, 20 L/fed., K-Humate, 10 L/fed., EM, 150 L/fed., compost tea + 10 L/fed., EM, 20 L/fed., K-Humate + 10 L/fed., EM and common chemical fertilizers (NPK). Amount of CO₂ emission were measured frequently every 30 days after sowing date (30, 60, 90, 120 and 150 days) and total accumulated CO₂. Moreover, Soil properties as available nitrogen (AN), soil organic matter (SOM), soil pH content and yield and its components were determined.

Results indicate that, the highest values of "AN" and "SOM" were recorded with compost tea + EM, while, the lowest values were obtained with chemical treatment. Chemical treatment increased soil pH more than other treatments except K-Humate and K-Humate + EM treatments. The compost tea + EM treatment gave the high reduction value of pH, while, control, K-Humate and K-Humate + EM treatments gave highest values without any significant between them. The greatest values of yield and its components were from applied compost tea + EM treatment. Meanwhile, the lowest values of tested parameters were recorded with added EM treatment. The lowest emission was taken by using compost tea + EM. While, the soil CO₂ emission was increased with added K-Humate + EM. In addition, amount of emission from soil was increased by the time to reach the peak after 120 days from sowing date. Furthermore, indicated that, the average air temperature, minimum air temperature, maximum air temperature and average soil temperature, respectively, had significant exponential relationship regression during the entire measuring period of CO₂ emission.

Keyword: Soil CO₂ emissions

1. INTRODUCTION

Carbon emissions related to human activities have been significantly contributing to the elevation of atmospheric (CO₂) and temperature. Recently, carbon emissions have greatly accelerated, thus much stronger effects on crops are expected (DaMatta *et al.*, 2009).

Carbon dioxide (CO₂) is an important greenhouse gas accounting for 60% of the total greenhouse effect. Soil is a major source for atmospheric CO₂. In the event of growing threats of global warming due to greenhouse gas emissions,



reducing CO₂ emission by isolate C in the soil is prime importance. Soil management practices like increasing soil organic carbon content, reduced tillage, manuring, residue incorporation, improving soil biodiversity, micro-aggregation and mulching can play an important role in isolate C in soil (Rastogi *et al.*, 2002).

The CO₂ emitted from soils is a useful indicator to determine energy flow patterns, specifically the mineralization of nutrients and the rate of organic material decomposition. The CO₂ flow from soils is the result of plant roots, organic matter mineralization, microflora, fauna, and to a small extent, by chemical oxidation of carbon-bearing materials. (Curtin *et al.*, 2000; Matteucci *et al.*, 2000; Al-Kaisi and Yin, 2005; Sainju *et al.*, 2008; Muñoz *et al.*, 2010).

The rate of soil CO₂ emission is normally controlled by several factors, such as CO₂ concentration gradient between the soil and the atmosphere, soil temperature, soil moisture, pore size, and wind speed. In addition, soil CO₂ emission is affected by agricultural practices such as irrigation, tillage, cropping system and N fertilization can alter crop residue C inputs, nutrient dynamics, soil temperature, residue management and varies with climatic conditions (Curtin *et al.*, 2000; Al-Kaisi and Yin, 2005; Sainju *et al.*, 2008). The measurement of soil CO₂ emission could provide a more sensitive indication of soil C sequestration than low-resolution data such as total or organic C values (Fortin *et al.*, 1996; Grant, 1997).

The atmospheric carbon dioxide (CO₂) concentration has increased about 85 ppm in the last 100 years (Lal, 2004) and approximately 10% of the CO₂ in the atmosphere passes through the soil each year (Wilson and Al-Kaisi; 2008). The emission of CO₂ was affected by the phenological stage of the plant while organic fertilizer increased the CO₂ emission (Fernández-Luqueño *et al.*, 2009).

Moreover, soil CO₂ emission is the main pathway of carbon emission from soil to atmosphere in terrestrial ecosystems and an important source of atmospheric CO₂ (Maljanen *et al.*, 2003; Koponen *et al.*, 2006; Nosalewicz *et al.*, 2013). Annual global soil respiration is estimated at 68–100 Pg year⁻¹ of C, accounting for almost 10% of the atmospheric CO₂ cycles and more than 11 times of the CO₂ released from fossil fuel combustion (Marland *et al.*, 2000). Therefore, knowledge of soil respiration dynamics and its controlling factors in different terrestrial ecosystems is essential to find proper management policies and relevant technologies to decrease soil CO₂ emissions and enhance carbon isolate.

Many studies have proved that organic fertilizer contributes to CO₂ emissions after soil application and the introduction of appropriate manure management techniques represents one opportunity for greenhouse gases mitigation (Bertora *et al.* 2008; Fernández-Luqueño *et al.* 2009; Cayuela *et al.* 2010).

Nitrogen has been regarded as a significant factor controlling soil respiration in N-deficient terrestrial ecosystems. Especially in the next few decades, with increasing rates of anthropogenic N deposition (Mosier *et al.*, 2002) and application of fertilizer, much N is to enter terrestrial ecosystems. This will change soil N status and plant N concentrations (Houghton, 2002). As a result, this will affect plant growth, microbial activities (Compton *et al.*, 2004; Frey *et al.*, 2004), litter decomposition and root respiration (Vose and Ryan, 2002), and thus soil respiration is expected to change. Many studies on the effect of N additions on soil respiration have been conducted (Bowden *et al.*, 2000; 2004; Burton *et al.*, 2004; Micks *et al.*, 2004) as well as agricultural ecosystems (Ding *et al.*, 2007; Al-Kaisi *et al.*, 2008).

Since the application of organic fertilizers became rare during the last decade, at present the main source of organic matter for cultivated soils are roots and stubble residues of agricultural plants. The rate of CO₂ emission from the soil is an important indicator of its microbial activity and intensity of organic matter decomposition. As a rule, the measurements of CO₂ emission rate are carried out through the growing period under field conditions and at a temperature above 5°C in the model experiments. It is suggested that CO₂ production and emission are very poor at low temperatures, and total CO₂ emission is negligible beyond the growing season (Lopes de Gerenyu *et al.*, 2005).

It is well known that CO₂ production, transport and emission in soil depend on environmental factors such as aeration condition, soil temperature, soil moisture, supplies of organic carbon, fertilization, pH, etc. (Weitz *et al.* 2001; Bowden *et al.* 2004; Lin *et al.* 2010).

Temperature (soil or air) is the best predictor of the annual and seasonal dynamics of CO₂ evolution rate of soils (Kirschbaum, 2000; Raich *et al.*, 2002; Perrin *et al.*, 2003). Meanwhile, soil temperature is the main ecological factors controlling the process of soil organic matter decomposition, CO₂ production and emission from soils (Rustad *et al.*,



2000). A high positive correlation between CO₂ emission rate and soil temperature was found for many soils under natural and agricultural conditions (Raich *et al.*, 2002; Lopes de Gerenyu *et al.*, 2005). In addition, a positive correlation between soil temperature and soil CO₂ efflux is well described by several reviews.

Concerning effects of air temperature, results of mathematical modeling suggest that, in mid- to high-latitude regions, moderate to medium local increases in temperature (1–3°C), along with associated CO₂ increase and rainfall changes, can have beneficial impacts on crop yields, but in low-latitude regions even moderate temperature increases (1–2 °C) are likely to have negative impacts on yield of major cereals (Easterling *et al.*, 2007).

The aims of the present study were first to quantify the seasonal fluxes of CO₂ from cultivated wheat with using different types of fertilizers. Second is to determine the impact of soil and air temperature on CO₂ evolution rate from the soil.

2. MATERIALS AND METHODS

Site and treatment description:

The present investigation was carried out on the experimental farm and laboratory of Soils and Water Sci. Dept., Fac. of Agric., AL-Azhar Univ., Assiut, Egypt, during two successive winter seasons of 2011/2012 and 2012/2013. The aim of conducted field experiment to study the effect of organic sources of nutrients use (Compost tea and K-Humate), and EM "Effective Microorganisms" bio-fertilization on soil CO₂ emissions as well as improving chemical and biological activity properties through 5 periods (30, 60, 90, 120 and 150 days from sowing date) and total accumulated CO₂. The experiment was designed as randomized completed blocks design with three replicates. Experimental treatments included six treatments as are presented as follow:-

1. NPK (Control).
2. 150 L/fed. Compost tea.
3. 20 L/fed. K-Humate.
4. 10 L/fed. EM.
5. 150 L/fed. Compost tea + 10 L/fed EM.
6. 20 L/fed. K-Humate + 10 L/fed EM.

Experimental plots area was 3 × 3.5 m = 10.5 m² (1/400 fed.). Physical and chemical properties of the experimental soil sample collected before planting wheat seeds were presented in Table (1):

Table 1: Soil physical and chemical properties of the soil.

Particle-Size distribution			Texture grade	Saturation capacity (%)	Field capacity (%)	Wilting point (%)	Bulk density (g/cm ³)	Particle density (g/cm ³)
Sand %	Silt %	Clay %						
Season 2011/12								
55.20	29.60	15.20	Silt loam	51.50	25.75	12.87	1.49	2.54
Season 2012/13								
57.68	27.31	15.01	Silt loam	52.32	26.16	13.08	1.50	2.57
C.E.C (cmolc kg ⁻¹)	CaCO ₃ (%)	O.M (%)	ECe (dSm ⁻¹)	pH soil past	Total-N (%)	Ava-N (mg/kg)	Ava-P (mg/kg)	Ava-K (mg/kg)
Season 2011/12								
13.90	1.29	1.59	1.43	7.12	0.18	62.72	11.57	169.41
Season 2012/13								
12.84	1.33	1.63	1.38	7.23	0.20	68.98	10.47	156.85



Sources of nutrients:

1. NPK: mineral fertilizer as ammonium nitrate, calcium super phosphate and potassium sulphate as they are commonly used for growing wheat plants and recommended by Ministry of Agriculture were applied to be the control treatment.
2. K-Humate: Potassium humate was extracted and using the classic alkali/acid fractionation procedure and analysis conducted according to standard methods (Valdrighi *et al.*, 1996).
3. Compost tea: Technical approaches to making are aerobic compost tea was procedure (Ingham, 2005), Chemical analysis of the compost tea and soil was conducted according to Page *et al.* (1982).
4. EM (Effective microorganisms): Technology of nature farming was introduced by Japanese scientists (Higa, 1989) was provided from the ministry of Agriculture.

The total amounts of K-Humate, compost tea, EM, K-Humate + EM and compost tea + EM add to soil during irrigation in equal three doses.

Table 2: Chemical analyses of organic fertilizers applied.

Organic fertilizer	pH (1:2.5)	EC dSm ⁻¹ (1:2.5)	O.M (%)	N (mgkg ⁻¹)	P (mgkg ⁻¹)	K (mgkg ⁻¹)
Season 2011/12						
K-Humate	11.92	4.49	14.82	961	48	1462
Compost tea	7.61	1.84	8.61	245	19	206
Season 2012/13						
K-Humate	12.62	4.69	14.51	954	50	1470
Compost tea	7.73	1.89	7.83	235	18	212

Cultivation and sampling:

Wheat seeds (*Triticum aestivum vulgar.*, CV. Sids 1) at rate of 51 kg/fed. were sown on December 1st under the surface irrigation method and harvested after 160 days from planting in the two seasons. Plant samples of one m² from each plot were harvested and ten plants were taken randomly to determine yield attributes (straw, grain and biological yield). Nitrogen was determined in wheat grain by the distillation in a Macro-Kjeldahle apparatus (Black, 1983) and protein content in wheat grains was determined according to (A. O. A. C., 2000).

Climatic data:

Climatic factors considered in this study were: maximum, minimum, average air temperature (°C) and soil temperature (°C). All climatic factors were measured and recorded by automatic weather station (Latitude 27.5 °N, Longitude 31.1°E and Altitude 71 m), Campbell scientific, Inc, Central Laboratory for Agricultural Climate (CLAC).

Soil sampling:

Surface soil samples (0-30 cm) were collected after wheat harvest to determine the available nitrogen "AN", soil organic matter content "SOM" and soil pH.

Soil CO₂ emission measurements:

Twenty five cm² plastic boxes were placed in each experimental plot with a Kilter glass bottle containing 100 ml of 1.0 normal sodium hydroxide placed in the middle of the plastic boxes to trap the evolved CO₂ (samples were taken at 60% soil moisture). The plastic boxes were sealed (air-tight) during all periods.

The glass bottles with 1.0 N sodium hydroxyl (NaOH) were retrieved daily at end period (30 days). At each retrieval time, the plastic boxes were left open up to half hour to prevent the development of anaerobic condition. The amount of CO₂-C trapped was determined by back titration according to Page *et al.* (1982).

The net CO₂-C was calculated by subtracting the value of the blank from that of the treatments, using the relationship of Anderson (2001).

$$CO_2 = (B - V) NE$$

Where:



B = Volume of HCl used to neutralize NaOH in the beaker to the end point of the blank.

V = Volume of acid used to neutralize NaOH in the beaker to the end point.

N = Normality of the acid.

E = Equivalent weight (to express data as C, E=6, while as CO = E= 22).

By the end of experiment, soil sample from each treatment was collected, air-dried, sieved through 2 mm sieve then kept for some chemical properties determination.

Statistical analysis:

Statistical analysis was performed by using the SAS system (SAS, 2005). Duncan’s multiple range tests was used for means separation. Significance was assessed at P < 0. 05.

3. RESULTS AND DISCUSSION

Climatic condition:

The air and soil temperature in the experimental site was shown in Figures (1 and 2). The mean monthly air temperature ranged from 13.45°C to 29.07°C and from 13. 93°C to 29.81°C, when soil temperature recorded from 14.03°C to 31.07°C and from 15.01°C to 32.03°C during the 2011/2012 and 2012/2013 wheat growing seasons, respectively. The mean monthly maximum air and soil temperatures during wheat growing season in 2011/2012 were slightly lower than that in 2012/2013. While, minimum air temperature in season 2011/2012 was slightly higher than that in 2012/2013, at all months except months of January. Moreover, soil minimum temperature was slightly higher at March, April and May in season 2012/2013 compare with season 2011/2012.




Figure 1: Average air temperature during two growing wheat seasons.






Figure 2: Average soil temperature during two growing wheat seasons.

Soil properties:

The soil properties (available nitrogen "AN", soil organic matter "SOM" and soil pH) were influenced significantly by different types of fertilizer. Generally, organic fertilizer and bio-fertilizer treatments enhanced "AN" and "SOM", while, reduced soil pH compared with chemical treatment.

Illustrated data in (Table. 3) obtained that the highest values of "AN" and "SOM" were recorded with Compost tea + EM treatment, when, the lowest values with chemical treatment. In addition, data in (Table. 3) reflected that the chemical treatment increased soil pH more than other treatments, except K-Humate and K-Humate + EM treatments. The Compost tea + EM treatment gave the high reduction value of soil pH, while, control, K-Humate and K-Humate + EM treatments gave highest values without any significant value between them. Those results were recorded during two growing seasons.

The result is supported by Ullah *et al.*, 2008 who reported that the organic matter content and availability of N in soil has been increased by organic fertilizer application. On the other hand, soil pH was increased with chemical application than organic application. Yadav *et al.*, 2002 and wells *et al.*, 2000, mentioned that the chemical properties of soil were influenced by different sources of soil nutrients (organic and chemical). Soil pH varied significantly with the treatments, it decreased with organic manures application and application of chemical + organic but increased with only chemical fertilizer application. Moreover, soil organic matter was decreased by chemical fertilizer application, but was increased with all types of organic manure application and that was recorded the highest with combined application. Whereas, Bouajila and Sanaa, 2011 reported that, add of compost or manure acted significantly on soil characteristics, especially the fertility and its productive capacity. Moreover, authors reported that improve soil organic level, resulting in a higher organic carbon content which contributes to diminish climatic heating.

Also, Constantin *et al.*, 2010 found a positive relationship between C and N inputs via cover crop biomass and soil C and N contents on three different soil types. Field experiments showed that plots with annual inputs of animal manure hold more soil C and N than plots receiving mineral fertilizers only (Christensen and Johnston, 1997). Hence, there are cumulative effects of cropping systems with abundant organic inputs on soil total N contents that could influence levels of plant-available N and thus yields of cereal crops. In contrast, residual effects of N added with mineral fertilizers are negligible (Petersen *et al.*, 2010). The result might be due to improvement of other physical and chemical properties for organic manure application compared to the chemical fertilizer application.

Table (3): Effect of different types of fertilization on soil properties available nitrogen, soil organic matter and soil pH during 2011/2012 and 2012/2013 seasons.

Treatments	First season			Second season		
	AN	SOM	pH	AN	SOM	pH
NPK	51.81 D	1.683 C	7.927 A	51.58 D	1.660 C	7.900 A
Compost tea	54.22 CD	2.090 B	7.780 C	54.03 CD	2.077 B	7.753 C
K-Humate	57.36 C	2.000 B	7.923 A	57.16 C	1.990 B	7.893 A
EM	57.57 C	2.016 B	7.847 B	57.35 C	2.007 B	7.817 B
Compost tea + EM	69.44 A	2.460 A	7.723 D	69.24 A	2.437 A	7.697 D
K-Humate + EM	64.92 B	2.063 B	7.927 A	64.73 B	2.053 B	7.897 A

Yield and yield components:

Data presented in (Table. 4) indicated that various studied treatments had a significant effect ($P < 0.05$) on straw, grain yield, biological yield, total content of nitrogen in grain (TCNG) and grain protein content (GPC). The greatest values of yield and its components were observed with applied compost tea + EM treatment. Meanwhile, the lowest values of tested parameters were recorded with added EM treatment. This results obtained at 2011/2012 and



2012/2013 seasons. These results are in a harmony with those reported by (Atta Allah and Mohamed, 2003; Sarwar, 2005; Yassen *et al.*, 2006; Javaid and Shah, 2010; Youssef, 2011).

Such satisfactory result might be due to the prevailing bio-fertilizer supplemented organic fertilizer improved soil properties, especially organic matter and N content. The application of bio-fertilizer in combination with organic or mineral fertilizers increased the efficiency of both organic and mineral N fertilizer, but apply the bio-fertilizer alone was ineffective in increasing yield. Thus, bio-fertilizers could be used as value-added soil amendments by supplementing organic and low chemical fertilizer rates for improving soil fertility and sustaining crop productivity (Abbasia and Yousra, 2012). Also, other authors mentioned the same results bio-fertilizer is enhancing soil biological activity, which improved nutrient mobilization from organic and chemical sources. Also, the bio-fertilizer plays a significant role in regulating the dynamics of organic matter decomposition and the availability of plant nutrients and increasing nitrogen fixer. In this case, Elsayed *et al.*, 2005; El-Garhi *et al.*, 2007; Badr *et al.*, 2009; Bahrani *et al.*, 2010; Abd El-Lattief, 2012 found positive effect on yield and yield attributes of wheat when inoculated with bio-fertilizer. Controlled field trials in Iran, Khavazi *et al.*, 2005, found that yield improvements of more than 20% have been observed for wheat as a result of application of bio-fertilizer inoculums.

Table (4): Effect of different types of fertilization on yield (straw, grain and biological yield) and its components (total content of nitrogen in grain and grain protein content) during 2011/2012 and 2012/2013 seasons.

Treatments	First season				
	Straw yield (ton/fed.)	Grain yield (ton/fed.)	Biological yield (gm/m ²)	TCNG (%)	GPC (%)
NPK	5.687 B	3.910 B	2271 B	2.423 BC	15.15 B
Compost tea	5.803 B	2.987 E	2082 BC	2.507 AB	15.67 AB
K-Humate	5.123 BC	3.700 C	2077 C	2.570 A	16.06 A
EM	4.947 C	2.897 E	1842 D	2.310 C	14.44 C
Compost tea + EM	6.543 A	4.347 A	2571 A	2.600 A	16.25 A
K-Humate + EM	5.793 B	3.377 D	2162 BC	2.441 B	15.25 B
Second season					
NPK	5.400 AB	3.718 B	2171 B	2.407 B	15.04 BC
Compost tea	5.509 AB	2.824 E	1984 B	2.487 AB	15.54 AB
K-Humate	4.829 B	3.507 C	1985 B	2.560 A	16.00 A
EM	4.661 B	2.703 E	1753 C	2.290 C	14.31 C
Compost tea + EM	6.231 A	4.158 A	2474 A	2.577 A	16.10 A
K-Humate + EM	5.499 AB	3.194 D	2070 B	2.407 B	15.04 BC

Soil CO₂ emission:

Soil CO₂ flux was fitted with applied organic and bio-fertilizations treatment. The results indicated that soil CO₂ emission was reduced with application of compost tea + EM treatment through determined period (60, 90, 120 and 150 days from sowing date), while, lowest emission after 30 days from sowing date was observed in the chemical treatment. While, the soil CO₂ emission was increased with added K-Humate + EM treatment at measurements period from 60 to 150 days from sowing date and with compost tea treatment after 30 days from sowing date (Table. 5). This results are recorded in both growing season 2011/2012 and 2012/2013. These results may be due to the influence of compost tea plus EM treatment on the soil properties (available nitrogen, soil organic matter and pH), Table (3), which lead to increase the "AN" and "SOM" and reduced soil pH. Some authors obtained that the application of farmyard manure induced a rapid increase in CO₂ emissions from the soil than CO₂ emissions from the soil that fertilized with mineral N (Jäger *et al.*, 2011).

Increasing N fertilizer application decreased soil CO₂ emissions (Wilson and Al-Kaisi, 2008). Kowalenko *et al.* (1978) and Fogg (1988) found increased N fertilization depressed soil CO₂ emissions. The effects of N deposition on soil



CO₂ emissions have also been studied (Micks *et al.*, 2004; Bowden *et al.*, 2004; Burton *et al.*, 2004). These studies reported results similar to Kowalenko *et al.*, 1978 and Fogg (1988). The reason for the depression of soil CO₂ emissions due to N fertilization is relatively unclear. Previous studies have suggested that the depression occurred because increased N reduced enzymatic activity (Bowden *et al.*, 2004; Burton *et al.*, 2004), decreased pH (Aerts and de Caluwe, 1999), or decreased microbial biomass (Soderstrom *et al.*, 1983).

Furthermore, several studies described a similar positive correlation of soil pH-value and soil CO₂ efflux (Andersson and Nilsson, 2001). Högberg *et al.* (2003) demonstrated the direct positive effect on soil emission with soil pH tolerance of the bacterial community. A biological activity of soil microorganisms is permitted between a soil pH of a minimum of 3 and a maximum of {7 to 8} (Scheffer and Schachtschabel, 2002). In addition, inside the mentioned values and otherwise constant conditions observed a nearly linear increase of soil CO₂ emission. Variation of the soil pH-value of the soil and between the single measurement points at each site showed a positive correlation with the CO₂ efflux. During simultaneous measurements with higher soil pH exhibited higher CO₂ fluxes (Reth *et al.*, 2005).

Meanwhile, soil CO₂ efflux influenced by other factors like, the root respiration (Kutsch *et al.*, 2001) and comprises total soil CO₂ efflux, and plants continuously excrete exudates into the soil. On other hand, due to application of bio-fertilizers the soil pH (KCl solution) raised CO₂ emission (Durinina *et al.*, 2001; Kutyova *et al.*, 2002).

Table (5): Effect of different types of fertilizers application on soil CO₂ emission (CO₂-C Kg/fed) during 2011/2012 and 2012/2013 seasons.

Treatments	First season					
	30	60	90	120	150	Accumulative
NPK	1.233 B	1.856 A	1.856 CD	2.002 AB	1.941 C	8.890 C
Compost tea	1.747 A	1.914 A	1.910 ABC	1.944 B	1.897 D	9.412 B
K-Humate	1.344 B	1.871 A	1.865 BCD	1.940 B	1.935 C	8.956 C
EM	1.655 A	1.900 A	1.915 AB	2.057 A	2.017 B	9.544 AB
Compost tea + EM	1.573 A	1.768 B	1.830 D	1.936 B	1.885 D	8.992 C
K-Humate + EM	1.714 A	1.934 A	1.932 A	2.073 A	2.043 A	9.694 A
	Second season					
NPK	1.239 C	1.868 AB	1.884 BC	2.016 AB	1.957 C	8.965 C
Compost tea	1.756 A	1.924 A	1.939 AB	1.955 B	1.910 D	9.483 B
K-Humate	1.353 BC	1.880 A	1.893 BC	1.951 B	1.947 C	9.024 C
EM	1.663 A	1.909 A	1.943 AB	2.069 A	2.030 B	9.615 AB
Compost tea + EM	1.580 AB	1.779 B	1.857 C	1.949 B	1.900 D	9.065 C
K-Humate + EM	1.727 A	1.953 A	1.959 A	2.084 A	2.054 A	9.777 A

Total accumulative CO₂ emission from soil:

Data in Table (5) noted that applied K-Humate + EM treatment gave the highest values of total accumulative soil CO₂ emission. While, the lowest values of total accumulative soil CO₂ emission were recorded with application of NPK, K-Humate and compost tea + EM treatments, respectively, without any significant differences between both treatments. In addition, obtained results from regression analyses confirmed that no significant effect on total accumulative CO₂ emission on wheat yield and yield components. These results are confirmed in both growing seasons 2011/2012 and 2012/2013.

Time effect on CO₂ emission measurements:

The amount of carbon dioxide emission from soil was different on particular period of measurements. The lowest rate of CO₂ emission evolution was observed on 30 days after sowing date. With the passage of time carbon dioxide emission increased, the highest rate of CO₂ was recorded after 120 and followed by 150 days from sowing date



without any significant differences between them (Table. 6). The mentioned trend was not harmony with (Włodarczyk, 2000; Rogalski *et al.*, 2008), whom reported that CO₂ emission decreased substantially after one day.

Table (6): time effect on soil CO₂ emission (CO₂-C Kg/fed.) during 2011/2012 and 2012/2013 seasons.

Time	30	60	90	120	150
First season	1.543 C	1.874 B	1.884 B	1.952 A	1.952 A
Second season	1.553 C	1.886 B	1.912 B	2.004 A	1.967 A

Relationships between soil CO₂ emission and air or soil temperature:

According to stepwise regression procedure indicated that, the average air temperature, minimum air temperature, maximum air temperature and average soil temperature, respectively, had significant exponential relationship regression during the entire measuring period of CO₂ emission. While maximum and minimum soil temperatures were not obtained any significant regression with CO₂ respiration in both growing seasons (Figures 3 and 4). On other hand, average soil temperature through period of CO₂ emission varied from 14.03 to 24.23°C and 16.58 to 25.6°C, respectively in 2011/2102 and 2012/2013. The relatively higher average soil temperature appeared in the months of growing season from February to April than other months. In months from December to April, soil temperature was always above 10°C in both seasons.

Similar trend was observed with minimum air temperature arranged between 2 to 8.87°C and 2.43 to 8.35°C, respectively in 2011/2102 and 2012/2013, and started for increasing from February to April months in first season and March to April months in second season. Minimum air temperature decreased was relatively low 10°C through all CO₂ emission months in two seasons.

Maximum air temperature recorded from 24.9 to 33.02°C and 27.76 to 39.87°C, respectively in 2011/2102 and 2012/2013. Maximum air temperature started rising from February to April in two growing seasons. Maximum air temperature increased above 30°C through all CO₂ emission months except December to February months in first season and January month in second season.

Average air temperature was ranged from 13.45 to 22.07°C and 13.93 to 22.07°C, in 2011/2102 and 2012/2013. Increasing average air temperature was started from February to April months. Average air temperature was always above 10°C through both seasons.

Our results showed that CO₂ flux was equally related with both air temperature (average, minimum and maximum) and average soil temperature at the time of CO₂ flux measurement.




Figure 3: Average air temperature through measurements period of CO₂ during two growing wheat seasons.




Figure 4: Average soil temperature through measurements period of CO₂ during two growing wheat seasons.

This could be attributed to the air temperature (average, minimum and maximum) were more correlated with CO₂ flux than soil temperature (maximum and minimum) (Parkin and Kaspar, 2003), air temperature influence soil temperature, which in turn, directly influence CO₂ flux from the soil surface (Bajracharya *et al.*, 2000a, b; Curtin *et al.*, 2000; Parkin and Kaspar, 2003). Parkin and Kaspar (2003) reported that CO₂ flux was better related with air temperature than with soil temperature because of plant residue at the soil surface. CO₂ flux was equally related with both soil temperature and daily average air temperature at the time of CO₂ flux measurement. The CO₂ flux was linearly related with soil temperature and daily average air temperature at the time of CO₂ measurement (Sainju *et al.*, 2008). Moreover, Soil warming can enhance the soil microbial activities and root growth sharply (Zheng *et al.*, 2009). This leads to an active decomposition of soil organic C and enhancement of plant-derived CO₂ release from root respiration (Fang *et al.*, 2009).

Prediction of soil CO₂ emission:

Furthermore, the most effective climatic factors could be used to predict amount of soil CO₂ emission ($R^2 = 0.629$, $P < .0001$) was selected (Average air temperature (T_{AV}), minimum air temperature (T_{AMin}), maximum air temperature (T_{AMax}) and average soil temperature (T_{SV}), respectively). Prediction equation for CO₂ emission from soil was concluded according to the mentioned statistical analysis and presented as follow:

$$Y = 0.713 + 0.26 T_{AV} + 0.23 T_{AMin} - 0.1 T_{AMax} + 0.06 T_{SV}$$

Where is: Y= soil CO₂ emission.

4. CONCLUSION

This study demonstrated how different types of fertilization affected on soil CO₂ emission (CO₂-C Kg/fed) from cultivated wheat yield. Statistical analysis showed that the soil properties were effect by the type of fertilizer. The "AN" and "SOM" represented the positive response for organic and bio-fertilizer compared with chemical fertilizer. In the same time, soil pH increased by applied chemical treatment more than other treatments except K-Humate and K-Humate + EM treatments. Moreover, the greatest values of yield (straw, grain yield, biological yield) and its components (total content of nitrogen in grain "TCNG" and grain protein content "GPC") were observed when applied compost tea + EM treatment. Meanwhile, the lowest values of tested parameters were recorded with added EM treatment. Soil CO₂ emission was reduced with compost tea + EM treatment through determents period (60, 90, 120 and 150 days from sowing date), when, lowest emission after 30 days from sowing was observed in the chemical treatment. While, the soil CO₂ respiration was increased with added K-Humate + EM treatment at measurements period from 60 to 150 days after sowing date and with compost tea treatment after 30 days after sowing date. In addition, the lowest rate of CO₂ evolution was observed on 30 days after sowing date. With the passage of time carbon dioxide emission increased, the highest rate of CO₂ was



recorded after 120 days from sowing date. The climatic factors as well as (average air temperature, minimum air temperature, maximum air temperature and average soil temperature) had significant exponential relationship regression during the entire measuring period of CO₂ emission. This statistical regression method can be used to predict soil CO₂ emission where soil CO₂ respiration is primarily dependent on climatic factors.

5. REFERENCES

- [1] Abbasia, M. K. and M. Yousra. 2012. Synergistic effects of biofertilizer with organic and chemical N sources in improving soil nutrient status and increasing growth and yield of wheat grown under greenhouse conditions. *Plant Biosystems*. 146(1): 181-189.
- [2] Abd El-Lattief, E. A. 2012. Improving bread wheat productivity and reduce use of mineral nitrogen by inoculation with *Azotobacter* and *Azospirillum* under arid environment in Upper Egypt. *International Conference on Applied Life Sciences (ICALS2012)*. 393-398.
- [3] Aerts, R. and H. de Caluwe. 1999. Nitrogen deposition effects on carbon dioxide and methane emissions from temperate peatland soils. *Oikos* 84-1: 44-54.
- [4] Al-Kaisi, M. M. and X. Yin. 2005. Tillage and crop residue effects on soil carbon and CO₂ emission in corn-soybean rotations. *J. Environ. Qual.* 34: 437-445.
- [5] Al-Kaisi, M. M., M. L. Kruse and J. E. Sawyer. 2008. Effect of nitrogen fertilizer application on growing season carbon dioxide emission in a corn-soybean rotation. *J. Environ. Qual.* 37: 325-332.
- [6] Andersson, S and S. I. Nilsson. 2001. Influence of pH and temperature on microbial activity, substrate availability of soil-solution bacteria and leaching of dissolved organic carbon in a mor humus. *Soil Biol. Biochem.* 33: 1181-1191.
- [7] A.O.A.C. 2000. Association of official agricultural chemists, official and tentative methods of analysis, 11th ed. Washington, D.C., USA.
- [8] Atta Allah, S. A. and G. A. Mohamed. 2003. Response of wheat grown in newly reclaimed sandy soil to poultry manure and nitrogen fertilization. *J. Agric. Sci., Mansoura Univ.* 28: 7531-7538.
- [9] Badr, E. A., O. M. Ibrahim and M. F. El-Kramany. 2009. Interaction effect of biological and organic fertilizers on yield and yield components of two wheat cultivars. *Egypt. J. Agron.* 31(1): 17-27.
- [10] Bahrani, A., J. Pourreza and M. Hagh Joo. 2010. Response of winter wheat to co-inoculation with *Azotobacter* and *Arbuscular Mycorrhizal* fungi (AMF) under different sources of nitrogen fertilizer. *American-Eurasian J. Agric. and Environ. Sci.* 8(1): 95-103.
- [11] Bajracharya, R. M., R. Lal and J. M. Kimble. 2000a. Erosion effects on carbon dioxide concentration and carbon flux from an Ohio alfisol. *Soil Sci. Soc. Am. J.* 64: 694-700.
- [12] Bajracharya, R. M., R. Lal and J. M. Kimble. 2000b. Diurnal and seasonal CO₂-C flux from soil as related to erosion phases in central Ohio. *Soil Sci. Soc. Am. J.* 64: 286-293.
- [13] Bertora, C., F. Alluvione, L. Zavattaro, J. W. Van Groenigen, G. Velthof and C. Grignani. 2008. Pig slurry treatment modifies slurry composition, N₂O, and CO₂ emissions after soil incorporation. *Soil Biol. Biochem.* 40: 1999-2006.



GLOBAL JOURNAL OF ADVANCED RESEARCH
(Scholarly Peer Review Publishing System)

- [14] Black, C. A. 1983. Methods of soil Analysis. Parts 1 and 2. Amer. Soc. Agron. Inc. Publ., Madison, Wisc., USA.
- [15] Bouajila, K. and M. Sanaa. 2011. Effects of organic amendments on soil physico-chemical and biological properties. *J. Mater. Environ. Sci.* 2: 485-490.
- [16] Bowden, R. D., G. Rullo and G. R. Stevens. 2000. Soil fluxes of carbon dioxide, nitrous oxide, and methane at a productive temperate deciduous forest. *J. Environ. Qual.* 29: 268–276.
- [17] Bowden, R. D., E. Davidson, K. Savage, C. Arabia and P. Steudler. 2004. Chronic nitrogen additions reduce total soil respiration and microbial respiration in temperate forest soils at the Harvard Forest. *For. Ecol. Manage.* 196: 43–56.
- [18] Burton, A. J., K. S. Pregitzer, J. N. Crawford, G. P. Zogg and D. R. Zak. 2004. Simulated chronic NO₃-deposition reduces soil respiration in Northern hardwood forests. *Glob. Ch. Biol.* 10: 1080–1091.
- [19] Cayuela, M. L., O. Oenema, P. J. Kuikman, R. R. Bakker and J. W. van Groenigen. 2010. Bioenergy by-products as soil amendments? Implications for carbon sequestration and greenhouse gas emissions. *GCB Bioenergy.* 2: 201–213.
- [20] Christensen, B. T. and A. E. Johnston. 1997. Soil organic matter and soil quality – lessons learned from long-term experiments at Askov and Rothamsted. *Dev. Soil Sci.* 25: 399–430.
- [21] Compton, J. E., L. S. Watrud and L. A. Porteous. 2004. Response of soil microbial biomass and community composition to chronic nitrogen additions at Harvard Forest. *For. Ecol. Manage.* 196: 143–158.
- [22] Constantin, J., B. Mary, F. Laurent, G. Aubrion, A. Fontaine, P. Kerveillant and N. Beaudoin. 2010. Effects of catch crops, no till and reduced nitrogen fertilization on nitrogen leaching and balance in three long-term experiments. *Agric. Ecosyst. Environ.* 135: 268–278.
- [23] Curtin, D., H. Wang, F. Selles, B. G. McConkey and C. A. Campbell. 2000. Tillage effects on carbon fluxes in continuous wheat and fallow-wheat rotations. *Soil Sci. Soc. Am. J.* 64: 2080-2086.
- [24] Durinina, E. P., E. C. Komarovskaya, T. U. Kuteva and I. A. Arhipchenko. 2001. Effect caused by biofertilizer Bamil on P tranformtion in Podzol soil and crop productivity. *Agrochemistry.* 11: 43-48.
- [25] DaMatta, F. M., A. Grandis, B. C. Arenque and M. S. Buckeridge. 2009. Impacts of climate changes on crop physiology and food quality. *Food Research International.* 43: 1814–1823.
- [26] Ding, W., Y. Cai, Z. Cai, K. Yagi and X. Zheng. 2007. Soil respiration under maize crops: effects of water, temperature, and nitrogen fertilization. *Soil Sci. Soc. Am. J.* 71: 944-951.
- [27] Easterling, W. E., P. K. Aggarwal, P. Batima, L. M. Brander, L. Erda and S. M. Howden. 2007. Food, fiber and forest products. In M. L. Parry, O. F. Canziani, J. P. Palutikof, P. J. van der Linden, and C. E. Hanson (eds.), *Climate change: Impacts, adaptation and vulnerability. Contribution of Working Group II to the fourth assessment report of the intergovernmental panel on climate change* (pp. 273–313). Cambridge: Cambridge University Press.
- [28] El-Garhi, A. S., N. A. Atia and S. E. E. Fouda. 2007. Effect of inoculating N-fixing bacteria (Cerealine) on wheat (*Triticum aestivum*, L.) growth and nutrient content. *Zagazig J. Agric. Res.* 34(2): 249-273.



GLOBAL JOURNAL OF ADVANCED RESEARCH
(Scholarly Peer Review Publishing System)

- [29] El-Sayed, M. Z., A. E. Abd El-Sattar, H. A. Basha and I. M. Abd El-Hammeed. 2005. Improvement of wheat productivity in newly reclaimed soil in Egypt. *Annals UMCS, Sec. E*:60: 113-121.
- [30] Fang, H. J., G. R. Yu, S. L. Cheng, T. H. Zhu, Y. S. Wang, J. H. Yan, M. Wang, M. Cao and M. Zhou. 2009. Effects of multiple environmental factors on CO₂ emission and CH₄ uptake from old-growth forest soils. *Biogeosciences*. 6: 1–13.
- [31] Fernández-Luqueño, F., V. Reyes-Varela, C. Martínez-Suárez, R. E. Reynoso-Keller, J. Méndez-Bautista, E. Ruiz-Romero, F. López-Valdez, M. L. Luna-Guido and L. Dendooven. 2009. Emission of CO₂ and N₂O from soil cultivated with common bean (*Phaseolus vulgaris* L.) fertilized with different N sources. *Science of the Total Environment*. 407: 4289–4296.
- [32] Fogg, K. 1988. The effect of added nitrogen on the rate of organic matter decomposition. *Biol. Rev.* 63: 433–472.
- [33] Fortin, M. C., P. Rochette and E. Pattey. 1996. Soil carbon dioxide fluxes from conventional and no-tillage small-grain cropping systems. *Soil Sci. Soc. Am. J.* 60: 1541–1547.
- [34] Frey, S. D., M. Knorr and J. Parrent. 2004. Chronic nitrogen enrichment affects the structure and function of the soil microbial community in a forest ecosystem. *For. Ecol. Manage.* 196: 159–171.
- [35] Grant, R. F. 1997. Changes in soil organic matter under different tillage and rotation: Mathematical modeling in ecosys. *Soil Sci. Soc. Am. J.* 61: 1159–1175.
- [36] Higa, T. 1989. Studies on application of microorganisms in nature farming. II: The practical application of effective microorganisms (EM). In: *Proceedings of 7th IFOAM Conference*, Ouagadougou, Burkina Faso, West Africa.
- [37] Högberg, M. N., E. Bååth, A. Nordgren, K. Arnebrant and P. Högberg. 2003. Contrasting effects of nitrogen availability on plant carbon supply to mycorrhizal fungi and saprotrophs – A hypothesis based on field observations in boreal forest. *New Phytol.* 160: 225–238.
- [38] Houghton, R. A. 2002. Terrestrial carbon sinks-uncertain explanations. *Biologist*. 49: 155–160.
- [39] Ingham, E. 2005. *The compost tea brewing manual*. US Printings, Soil Foodweb Incorporated, Oregon.
- [40] Jäger, N., C. F. Stange, B. Ludwig and H. Flessa. 2011. Emission rates of N₂O and CO₂ from soils with different organic matter content from three long-term fertilization experiments—a laboratory study. *Biol. Fertil. Soils*. 47: 483–494.
- [41] Javaid, A. and M. B. M. Shah. 2010. Growth and yield response of wheat to EM (effective microorganisms) and parthenium green manure. *Afr. J. Biotechnol.* 9: 3373-3381.
- [42] Khavazi, K., H. Asadi-Rahmani and M. J. Malakouti. 2005. Necessity for the production of biofertilizers in Iran (eds.). Ministry of Jihad-e- Agriculture, Agricultural Research and Education Organization (AREO) and Soil and Water Research Institute (SWRI), Tehran, Iran, pp: 419.
- [43] Kirschbaum, M. U. F. 2000. Will changes in soil organic carbon act as a positive or negative feedback on global warming. *Biogeochemistry*. 48: 21–51.



GLOBAL JOURNAL OF ADVANCED RESEARCH
(Scholarly Peer Review Publishing System)

- [44] Koponen, H. T., C. E. Duran, M. Maljanen, J. Hytonen and P. J. Martikainen. 2006. Temperature responses of NO and N₂O emissions from boreal organic soil. *Soil Biology and Biochemistry*. 38: 1779-1787.
- [45] Kowalenko, C. G., K. S. Ivarson and D. R. Cameron. 1978. Effect of moisture content, temperature and nitrogen fertilization on carbon dioxide evolution from field soils. *Soil Biology and Biochemistry*. 10: 417-423.
- [46] Kutsch, W. L., A. Staack, J. Wotzel, U. Middelhoff and L. Kappen. 2001. Field measurements of root respiration and total soil respiration in an alder forest. *New Phytol*. 150: 157-168.
- [47] Kutuyova, T. Y., E. P. Durinina, N. E. Muravyova and A. V. Sheyko. 2002. Microbial fertilizers Bamil, Omug, Ekud, Pudret their properties, influence on soil and crops. *Herald of Moscow State University, Soil Science series*. 4: 40-46.
- [48] Lal, R. 2004. Soil carbon sequestration impacts on global climate change and food security. *Science*. 304: 1623-1627.
- [49] Lin, B., L. Chambers, P. Stackhouse Jr., B. Wielicki, Y. Hu, P. Minnis, N. Loeb, W. Sun, G. Potter, Q. Min, G. Schuster and T. F. Fan. 2010. Estimations of climate sensitivity based on top-of-atmosphere radiation imbalance. *Atmos. Chem. Phys*. 10: 1923-1930.
- [50] Lopes de Gerenyu, V. O., I. N. Kurganova, L. N. Rozanova and V. N. Kudeyarov. 2005. Effect of soil temperature and moisture on CO₂ evolution rate of cultivated Phaeozem: analysis of a long-term field experiment. *Plant Soil Environ*. 51(5): 213-219.
- [51] Maljanen, M., A. Liikanen, J. Silvola and P. J. Martikainen. 2003. Measuring N₂O emissions from organic soils with closed chamber or gas gradient methods. *Eur. J. Soil Sci*. 54: 625-631.
- [52] Marland, G., T. A. Boden and R. J. Andres. 2000. Global, regional and national fossil fuel CO₂ emissions. In: *Trends: a compendium of date on global change*. Oak Ridge National Laboratory, U.S. Oak Ridge, Tennessee, USA.
- [53] Matteucci, G., S. Dore, C. Rebmann, S. Stivanello and N. Buchmann. 2000. Soil respiration in beech and spruce forest in Europe: trends, controlling factors, annual budgets and implications for the ecosystem carbon balance. p. 217-236. In E.D. Schulze (ed.) *Carbon and nitrogen cycling in European forest ecosystems*. Springer Verlag, Berlin, Germany.
- [54] Micks, P., J. D. Aber, R. D. Boone and E. A. Davidson. 2004. Short-term soil respiration and nitrogen immobilization response to nitrogen applications in control and nitrogen-enriched temperate forests. *For. Ecol. Manage*. 196: 57-70.
- [55] Mosier, A. R., J. A. Morgan and J. Y. King. 2002. Soil-atmosphere exchange of CH₄, CO₂, NO and N₂O in the Colorado Short grass Steppe under Elevated CO₂. *Plant Soil*. 240: 201-211.
- [56] Muñozl, C., L. Paulino, C. Monreal and E. Zagal. 2010. Greenhouse gas (CO₂ and N₂O) emission from soils: a review. *CHIL. J. AGR. RES*. 70(3): 485-497.
- [57] Nosalewicz, M., Z. Stêpniewska and A. Nosalewicz. 2013. Effect of soil moisture and temperature on N₂O and CO₂ concentrations in soil irrigated with purified wastewater. *Int. Agrophys*. 27: 299-304.
- [58] Page, A. L., R. H. Miller and D. R. Keeney. 1982. *Methods of Soil Analysis*. 2nd edn., Amercen Society of Agronomy, Madison, WI., USA.



GLOBAL JOURNAL OF ADVANCED RESEARCH
(Scholarly Peer Review Publishing System)

- [59] Parkin, T. B. and T. C. Kaspar. 2003. Temperature controls on diurnal carbon dioxide flux: Implications for estimating soil carbon loss. *Soil Sci. Soc. Am. J.* 67: 1763-1772.
- [60] Perrin, D., E. Laitat, M. Yernaux, Q. Mezoesy and M. Aubinet. 2003. Temporal and spatial changes in the soil CO₂ efflux in a mixed temperate forest (Vielsalm, Belgium). *Comparative biochemistry and Physiology: SEB Abstracts*. 134, 3: 191.
- [61] Petersen, J., I. K. Thomsen, L. Mattsson, E. M. Hansen and B. T. Christensen. 2010. Grain yield and crop N off take in response to residual fertilizer N in long-term field experiments. *Soil Use Manage.* 26: 455-464.
- [62] Raich J. W., C. S. Potter and D. Bhagavatti. 2002. Interannual variability in global soil respiration, 1980-94. *Global Change Biol.* 8: 800-812.
- [63] Rastogi, N. K., K. S. M. S. Raghavarao, K. Niranjana and D. Knorr. 2002. Recent developments in osmotic dehydration: Methods to enhance mass transfer. *Trends in Food Science and Technology*. 13: 48-59.
- [64] Reth, S., M. Reichstein and E. Falge. 2005. The effect of soil water content, soil temperature, soil pH-value and the root mass on soil CO₂ efflux – A modified model. *Plant and Soil*. 268: 21-33.
- [65] Rogalski, L., A. Beś and K. Warmański. 2008. Carbon dioxide Emission to the atmosphere from overburden under controlled temperature conditions. *Polish J. of Environ. Stud.* 17(3): 427-432.
- [66] Rustad, L. E., T. G. Huntington and R. D. Boone. 2000. Controls on soil respiration: Implication for climate change. *Biogeochemistry*. 48: 1-6.
- [67] Sainju, U. M., J. D. Jabro and W. B. Stevens. 2008. Soil carbon dioxide emission and carbon sequestration as influenced by irrigation, tillage, cropping system and nitrogen fertilization. *J. Environ. Qual.* 37: 98-106.
- [68] Sarwar, G. 2005. Use of compost for crop production in Pakistan. Ph.D. Dissertation (Published) *Ökologie und Umweltsicherung*. Universität Kassel, Fachgebiet Landschaftsökologie und Naturschutz, Witzenhausen, Germany.
- [69] SAS Institute. 2005. The SAS system for Microsoft Windows. Release 9. 1. SAS Inst., Cary, NC.
- [70] Scheffer, F. and P. Schachtschabel. 2002. *Lehrbuch der Bodenkunde*. Spektrum, Akad. Verl., Heidelberg, Germany, 593 p.
- [71] Soderstrom, B., E. Baath and B. Lundgren. 1983. Decrease in soil microbial activity and biomasses owing to nitrogen amendments. *Can. J. Microbiol.* 29: 1500-1506.
- [72] Ullah, M. S., M. S. Islam, M. A. Islam and T. Haque. 2008. Effects of organic manures and chemical fertilizers on the yield of brinjal and soil properties. *J. Bangladesh Agril. Univ.* 6(2): 271-276.
- [73] Valdrighi, M. M., A. Pera, M. Agnolucci, S. Frassinetti, D. Lunardi and G. Vallini. 1996. Effects of compost-derived humic acids on vegetable biomass production and microbial growth within a plant (*Cichorium intybus*)-soil system: a comparative study. *Agriculture, Ecosystems and Environment*. 58: 133-144.
- [74] Vose, J. M. and M. G. Ryan. 2002. Seasonal respiration of foliage, fine roots, and woody tissues in relation to growth, tissue N, and photosynthesis. *Global Change Biol.* 8: 182-193.



GLOBAL JOURNAL OF ADVANCED RESEARCH
(Scholarly Peer Review Publishing System)

- [75] Weitz, A. M., E. Linder, S. Frolking, P. M. Crill, and M. Keller. 2001. N₂O emissions from humid agricultural soils: effects of soil moisture, texture and nitrogen availability. *Soil and Biogeochemistry*. 33: 1077–1093.
- [76] Wells, A. T., K. Y. Chan and P. S. Cornish. 2000. Comparison of conventional and alternative vegetable farming system on the properties of a yellow earth in New South Wales. *Agriculture, Ecosystems and Environment*. 80 (1/2): 962-966.
- [77] Wilson, H. M. and M. M. Al-Kaisi. 2008. Crop rotation and nitrogen fertilization effect on soil CO₂ emissions in central Iowa. *Applied Soil Ecology*. 39: 264-270.
- [78] Włodarczyk, T. 2000. N₂O emission and absorption against a background of CO₂ in eutric cambisol under different oxidation-reduction control. *Acta Agrophisica* 28: 39-43.
- [79] Yadav, A. C., S. K. Sharma and B. R. Batra. 2002. Effect of sodic water, FYM and gypsum on the soil, growth and yield of brinjal. *Ann. Agric. Biol. Res.* 7(1): 73-77.
- [80] Yassen, A. A., M. Abd El-Hady and S. M. Zaghoul. 2006. Replacement part of mineral N fertilizer by organic ones and its effect on wheat plant under water regime conditions. *World J. Agric. Sci.* 2: 421-428.
- [81] Youssef, M. A. 2011. Synergistic impact of effective microorganisms and organic manures on growth and yield of wheat and marjoram plants. Ph. D. Fac. Agric., Assiut Univ., Egypt.
- [82] Zheng, Z. M., G. R. Yu, Y. L. Fu, Y. S. Wang, X. M. Sun and Y. H. Wang. 2009. Temperature sensitivity of soil respiration is affected by prevailing climatic conditions and soil organic carbon content: A trans-China based case study. *Soil Biology and Biochemistry*. 41: 1531–1540.